

University of Groningen

Symmetry breaking in navigating cells

Kataria, Rama

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2015

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kataria, R. (2015). *Symmetry breaking in navigating cells: The role of heterotrimeric and monomeric G-proteins in Dictyostelium chemotaxis*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Chapter I

Introduction

Rama Kataria, Peter J.M. van Haastert, Arjan Kortholt

Department of Cell Biochemistry, University of Groningen, Nijenborgh 7, 9747 AG, Groningen,
The Netherlands

INTRODUCTION

Cells have the remarkable property to sense and migrate towards extracellular gradients of chemoattractants. This directional movement of cells, also called chemotaxis, is important for almost all organisms. Prokaryotes rely on chemotaxis to sense nutrients, toxins, oxygen levels, pH and osmolarity, while metazoans need chemotaxis for embryonic organization [1], tissue morphogenesis [2] and the immune response by leukocytes [3]. Defects in chemotaxis are associated with the development of various diseases such as cancer [4,5], asthma, arthritis, inflammatory bowel disease and cardiovascular diseases [6]. Therefore, it is important to elucidate the exact mechanisms regulating directional migration of cells.

***Dictyostelium* as a model organism**


The most commonly used model systems for studying eukaryotic chemotaxis are the slime mold *Dictyostelium discoideum* and neutrophils. Although these model systems have clearly distinct roles, they have a highly similar chemotactic behavior [7]. They display strong chemotactic responses, their stimuli are well-defined and they are highly sensitive to even shallow gradients (~ 2% across the length of their body) of chemoattractants. *Dictyostelium* has a remarkable life cycle, consisting of a unicellular and multicellular phase. During the vegetative stage, single cells divide by mitosis as they feed themselves on bacteria. The bacteria secrete folic acid, which is a chemoattractant for the *Dictyostelium* cells. Upon starvation, *Dictyostelium* cells undergo a drastic change in gene expression and enter the multicellular phase of its life cycle, which is characterized by development and cell differentiation. The cells start to secrete cAMP and simultaneously sense cAMP secreted by the neighbouring cells thus forming a gradient of cAMP around the origin point. cAMP synthesis and secretion by *Dictyostelium* cells is an oscillatory process. The outward propagation of cAMP is regulated by extracellular and intracellular phosphodiesterases (PDEs) that degrade the accumulated cAMP. This degradation of cAMP by PDEs serves to maintain the steepness of cAMP gradient thereby limiting the interval and degree of signaling. As a result cAMP waves from the initiation centre are propagated every ~ 6 min. These synchronized oscillatory waves of cAMP results in the formation of streams moving along the gradient towards the centre. After 6 hours aggregates are formed and cells start to differentiate between prestalk cells, prespore cells and anterior-like cells (ALC) depending on cell cycle position and state of nutrition. The tipped mound develops into slugs after 16 hours of

starvation. The anterior region of the slug is occupied by prestalk cells, while prespore cells intermingled with the anterior-like cells are present at the posterior end. Settlement of the slug and the subsequent formation of a Mexican hat, mark the beginning of the culmination stage. The prestalk and prespore cells switch positions at the time of culmination to form a fruiting body comprising of a stalk of dead vacuolated prestalk cells and a spore head called sorus on top made by prespore cells (Fig 1). *Dictyostelium* spores can survive harsh unfavourable conditions. Under better conditions the spores germinate to release single celled amoeba in the vegetative stage, thus completing the life cycle.

We use *Dictyostelium* as model system for studying chemotaxis and development because it is a haploid organism with easy genetics, it has a short and simple life cycle, can easily be grown in large quantity for biochemical experiment, and its complete 34Mb genome is sequenced and available on dictybase [8,9]. Most importantly many molecular features regulating chemotaxis are conserved between *Dictyostelium* and higher eukaryotes [7,10].

Chemotaxis

Chemotaxis is a complex cellular process involving three separate but inter-related events: directional sensing, cellular polarity and cell motility [11]. Cell detects an asymmetric extracellular gradient by sensing the changes in receptor occupancy. Previous studies have shown that *Dictyostelium* can sense the receptor occupancy as a function of space (spatial sensing) and time (temporal sensing) [11]. The shallow extracellular gradient of a chemoattractant is transduced into a steep gradient of intracellular signaling components resulting in selective localization of the signaling molecules to one end of a cell facing the gradient. This process also referred to as symmetry breaking [12,13], induces cytoskeletal rearrangement with F-actin polymerization and pseudopod extension at the side of the cell facing the gradient (leading edge) and myosin II assembly at the back of the cell (lagging edge) [14-16]. Pseudopod formation is biased at the leading edge as it is more sensitive to gradients, while lateral pseudopod formation is suppressed. This results in faster and more directional movement of cell towards the direction of chemoattractant. Cells treated with inhibitors of actin polymerization become round and are incapable to move, but still can sense the direction of the gradient and show symmetry breaking [13,17]. Thus directional sensing and symmetry breaking are independent of cell elongation or cell motility [12].



Chisholm, R.L. et al., Nat. Rev. Mol. Cell Biol. 5, 531–541 (2004).

Figure 1 Life cycle of *Dictyostelium discoideum* During vegetative phase, individual *Dictyostelium* cells feed on soil living bacteria. Upon starvation, cells undergo a drastic change in gene expression and enter the developmental phase where each cell begins to secrete cAMP and chemotax towards cAMP secreted by other cells. This leads to formation of aggregates which differentiate to form slug-like multicellular body which in turn, culminates to form a fruiting body consisting of stalk of vacuolated dead cells and spore head on top.

Gradient detection by GPCR

The initial event in directional sensing of eukaryotic cells is the binding of chemoattractant to G-Protein Coupled Receptors (GPCRs) that are uniformly distributed over the cell surface. GPCRs are ubiquitously present in all eukaryotes with humans having more than 800 GPCRs grouped in five families [18]. The *Dictyostelium* genome encodes for more than 30 GPCRs, including the cAMP receptors, and receptors belonging to GABA, PAF and serotonin families [19]. GPCRs are characterized by seven membrane spanning α -helices with the N-terminal extracellular domain acting as a ligand-binding domain for large ligands such as peptide hormones and small ligands such as cAMP bind in the barrel formed by the seven α -helices. *Dictyostelium* contains four cAMP receptors (cAR1-CAR4) [20,21], that show 54-69% amino acid identity and are

expressed during different stages of the life cycle [22]. cAR1 has the highest affinity for cAMP and it is expressed during early developmental stages. Cells lacking *car1* are insensitive to cAMP, do not show any chemotaxis and fail to initiate development [23]. The expression of cAR1 is low during growth, goes up during early starvation, and decreases again at later stages of development [24]. cAR2 is majorly expressed after 15h of development in anterior prestalk cells. *car2*-null cells do not form tips and development arrests after the tight aggregate stage [25,26]. On the other hand, *car3*-null cells display no characteristic phenotype; they aggregate and develop normally [27]. The level of cAR3 is peaking at mound stage, after 10h of development, and becomes restricted to prespore cells [25]. cAR4 expression starts after tip elongation and drops after the culmination stage. Only 25% of *car4*-null cells are able to form very short fruiting bodies [20]. Although much is known about folate chemotaxis, the folate receptor, through which vegetative *Dictyostelium* cells tracks bacteria has still not been identified.

Heterotrimeric G-proteins

In the absence of ligand, GPCRs are coupled to an inactive heterotrimeric complex consisting of a GDP bound $G\alpha$ -subunit and a $G\beta\gamma$ dimer. Upon ligand binding, the GPCR gets activated and promotes the dissociation of GDP from the $G\alpha$ -subunit. The nucleotide-free $G\alpha$ subunit then rapidly binds GTP which is present in higher concentration in the cytoplasm. Thus GPCRs function as GEFs (Guanine nucleotide Exchange Factors) stimulating the exchange of GDP to GTP on $G\alpha$ subunit [28-30]. $G\alpha$ -GTP undergoes a conformational change and dissociates from $G\beta\gamma$, so that both subunits are free to activate downstream signal transduction pathways. The signal is terminated by the intrinsic GTPase activity of $G\alpha$ -subunits that mediates the hydrolysis of GTP to GDP. RGS proteins (Regulators of G-protein Signaling) can enhance the intrinsic GTPase activity and thereby accelerate the hydrolysis reaction [31,32]. The $G\alpha$ -GDP subunit reassociates with $G\beta\gamma$ to form the inactive heterotrimer again [33,34] (Fig 2A).

Dictyostelium contains only a single $G\beta$ [35] and $G\gamma$ [36] protein. *Dictyostelium* $G\beta$ shares 60% homology with that of human, *Drosophila* and *C.elegans* [35]. The C-terminal region of $G\beta$ consists of seven WD40 repeats that fold to form a β -propeller structure, which serves as binding site for $G\alpha$ -subunits and effectors [34,37,38]. The N-terminal region forms a tight coiled-coil structure with $G\gamma$. *g\beta*-null cells are viable but fail to aggregate upon starvation

[35]. These cells also lack the ability to chemotax towards folate or cAMP and are defective in cAMP-stimulated activation of guanylyl cyclase and adenylyl cyclase [38]. Consistently, over-expression of a mutant of G γ , in which four conserved residues (CSVL) at C-terminal have been deleted, results in decreased membrane localization of G $\beta\gamma$, and subsequently failure of aggregation [36]. This shows that G β is essential for almost all chemoattractant-mediated responses and plays a central role in transducing extracellular signals.

The *Dictyostelium* genome encodes for twelve Ga subunits [39], which are most similar to the mammalian Gai family [39]. They have distinct roles during the life cycle of *Dictyostelium* [40-45] (Table1). Ga2 and Ga4 are essential for chemotaxis towards cAMP and folate, respectively. Ga2 binds to cAR1 and is essential for all cAMP mediated responses [41,46]. Ga4 is expressed during vegetative growth and at low level during early development; the expression increases during mound stage and remains high throughout the development [47]. The subunit Ga4 is essential for folate chemotaxis and during late development stages for the formation of anterior prestalk cells and spores [47,48].

The heterotrimeric G-proteins, are like the GPCRs, localized uniformly on the plasma membrane [49,50]. This ensures the detection of chemoattractant over the whole cell body [49,51]. Previous studies have shown that receptor occupancy and heterotrimeric G-protein activation reflects the steepness of the extracellular gradient of chemoattractant [52]. This suggest that amplification of the extracellular signal and symmetry breaking occurs first downstream of receptor and heterotrimeric G-protein activation [53,54] (Fig 2B).

Small G-proteins

Ras family

One of the first downstream responses of heterotrimeric G-protein signaling is the activation of Ras proteins [55,56]. Ras belongs to the family of small G-proteins, also known as small GTPases that act as molecular switches. G-proteins cycle between an inactive GDP-bound and active GTP-bound state. This cycle is tightly regulated by GEFs which catalyze the exchange of GTP for GDP, thereby activating the Ras protein. GAPs convert the Ras protein back into the inactive GDP-bound form by stimulating the otherwise low intrinsic hydrolysis activity by many orders of magnitude (Fig 2A). Like heterotrimeric G-proteins, Ras proteins are also uniformly




Figure 2 The G-protein cycle (A) Both heterotrimeric and monomeric G-proteins switch between the inactive GDP-bound state and the active GTP-bound state. For heterotrimeric G-proteins, GPCRs (G-protein coupled receptors) act as GEFs (Guanine nucleotide exchange factors) catalyzing the exchange of GDP to GTP. The RGS proteins (Regulators of G-protein cycle) assist the intrinsic GTPase activity of $G\alpha$ proteins to hydrolyze the bound-GTP and return to the inactive state. RasGEFs and RasGAPs assist the Ras cycle in a similar fashion. The G-proteins interact with the downstream effectors in the active GTP-bound state. (B) Receptor occupancy and the activation of associated heterotrimeric G-proteins are proportional to the steepness of extracellular chemoattractant gradient; which differs 2% across the length of the cell. Due to internal amplification, the activation of Ras is much stronger at the front as compared to the rear of the cell.

Table 1 Gα- proteins in *Dictyostelium*

Gα-proteins in <i>Dictyostelium</i>	Expression profile	Function	Ref
Gα1	expressed predominantly in anterior prestalk cells, expression becomes higher during mound stage and continues till culmination stage.	controls prestalk morphogenesis, regulates cGMP pathway, OE cells are large & multinucleate.	[57,58] [59]
Gα2	expressed at low levels during growth & induced to high levels during aggregation.	cAMP chemotaxis & relay, activation of adenyly and guanylyl cyclase.	[46,58] [59]
Gα3	expression induced to high level at 2 h of starvation, then gradually declines	cAMP production, inhibits PKA activity	[44,60] [61]
Gα4	expressed during growth, getting higher at mound stage n continuing till culmination.	folate chemotaxis, late development	[45,47] [62]
Gα5	same as temporal expression of Gα4	inhibit folate chemotaxis, tip morphogenesis	[42,63] [64]
Gα6	expressed during growth and early development		
Gα7	expressed during aggregation and mound stage	Gα7 OE form abnormal fruiting bodies	[65]
Gα8	expressed at low level during growth, reaching maximum at aggregation and mound stage	cell proliferation & differentiation cell-substrate adhesion, proliferation inhibiting activity of AprA.	[65,66] [67,68]
Gα9	expressed throughout growth & development, reach maximum after 5 hours of starvation	delays the expression of cAMP regulated genes & inhibits the formation of cAMP initiation centres during early development.	[39]

distributed along the plasma membrane and are exclusively localized at the leading edge in a chemotaxing cell [56,69]. Contrary to activation of $G\alpha\beta\gamma$ which is proportional to the steepness of gradient, activation of Ras is much stronger as a result of internal amplification (Fig 2B). For such a simple organism, *Dictyostelium* contains an unusually high amount of Ras subfamily proteins; it has 11 Ras proteins, 3 Rap proteins (RapA, B, and C) and one Rheb protein [69,70]. Ras subfamily proteins are involved in many important and diverse processes like growth, chemotaxis, development, cytokinesis and regulation of cytoskeleton (Table 2).

Table 2 Ras proteins in *Dictyostelium*

Protein	Expression profile	Function	Ref
RasB	expressed predominantly during growth and first 8 hours of development	regulation of cell cycle	[71,72]
RasC	expressed throughout life cycle	cAMP relay, development, activation of ACA and PKB	[73,74] [75,76]
RasD	expressed during growth, reappear in tight aggregate stage	phototaxis, thermotaxis cAMP production	[77,78]
RasG	expressed during growth and early development	growth, chemotaxis & development	[76,79] [80,81]
RasS	expressed during starvation and aggregation	regulates the balance between eating and moving	[82,83] [84]
RapA	expressed at all stages but maximally during aggregation and culmination	growth, chemotaxis and cytoskeleton rearrangement	[85,86] [87,88]

RasC and RasG are the best characterized small G-proteins and together important for chemotaxis and development [55,76,81]. RasC is important for proper adenylyl cyclase and PKB activation, which is essential for cAMP relay and development [73,76]. *rasC*-null cells have reduced chemotaxis and abolished aggregation. Cells lacking *rasG* have growth defect, strongly reduced chemotaxis, delayed aggregation and the aggregate size is smaller than wild type cells. cAMP induced accumulation of cGMP and phosphorylation of PKB is also reduced in *rasG*-null cells [80],[81]. Disruption of both *rasC* and *rasG* leads to a complete loss of cAMP relay and

chemotaxis, suggesting that RasC and RasG are together responsible for activating all major downstream pathways [81].

Regulation of the Ras cycle

Recent studies have given important insight in the network regulating Ras activity; however the exact pathways remain largely unknown. In the *Dictyostelium* genome, there are 25 genes encoding RasGEFs and 17 genes for putative RasGAPs [89]. Disruption of the GAP, *nfaA*, results in severe chemotaxis defects, delayed aggregation, increased levels of Ras activation and F-actin. NfaA regulates both RasB and RasG [90]. RasGEFA is essential for the activation of RasC [55,76]. *gefA*⁻ cells are completely lacking RasC activation, and consistently have normal growth and chemotaxis, whereas activation of ACA and development are severely impaired [74,81,91]. RasGEFR specifically activates RasG *in vitro* [55]. Although, *gefR*⁻ cells have reduced cAMP mediated RasG activation, there is still a significant level of active RasG, suggesting there have to be additional RasG specific GEFs.

Since Ras signaling is essential for chemotaxis, cAMP relay and development, it will be essential to further unravel the complex network of GEFs and GAPs that regulate the spatial and temporal activation of Ras.

Developmental pathways

Upon starvation, the aggregation of *Dictyostelium* cells is driven by the synthesis and secretion of cAMP in an oscillatory manner and the ability of cells to sense and respond to it. This is followed by signal relay which involves the activation of Adenylyl Cyclase A (ACA) and subsequent secretion of cAMP [92] which is mainly regulated by RasC via activation of TORC2 pathway [93,94]. Activated TORC2 phosphorylates PKBR1 and PKBA which together play a critical role in signal relay by stimulating ACA [93] (Fig 3). TORC2 is a multi-subunit complex comprising of TOR kinase, PIA, RIP3 and Lst8 [95]. The *rip3*⁻, *pia*⁻ and *lst8*⁻ cells fail to aggregate upon starvation and accumulate less ACA [95-97], thereby, strengthening the role of TORC2 complex in development [95].

Dictyostelium genome encodes only two extracellular-signal related kinases sharing 37% homology: ERK1 and ERK2, which are expressed during growth and throughout development [98-100]. ERKs are involved in many vital cellular processes including cell proliferation, gene




Figure 3 Signaling pathways regulating development The activation of heterotrimeric G-proteins by binding of cAMP to cAR1 leads to activation of Ras proteins and MAP kinase pathways. The activation of RasG, RasC and MEKA leads to recruitment of PI3K, TorC2 and ERK respectively, which is important for activation of adenylyl cyclase A (ACA) and hence cAMP relay. ERK2 inhibits the cAMP specific phosphodiesterases RegA leading to increased levels of cAMP important for cell aggregation.

expression, growth, differentiation and apoptosis [101]. In other organisms, signal transduction to ERKs is considered to be regulated by Raf kinases which are activated by Ras proteins via a kinase cascade consisting of MAPK/MEK/ERK in mammalian system [101-103]. However recent studies revealed a Ras-independent mechanism for ERK activation [104,105]. *Dictyostelium* ERKs are a part of kinase cascade that includes MEKA and MKKA. MEKA is required for normal chemotaxis to cAMP, accumulation of guanylyl cyclase in response to cAMP and for aggregation [106]. MKKA regulates morphogenesis, developmental timing and cell-specific patterning during multicellular development [107]. *erk1*-null cells form small aggregates which may or may not form tips and the ones which succeed to form tips and

complete the development cycle have smaller fruiting bodies. ERK2 play a minor role in chemotaxis to cAMP and folate and is also involved in cell differentiation and development of prespore cells [108,109]. Moreover, ERK2 is also known to regulate cAMP levels through RegA, a cAMP-specific phosphodiesterases [110] (Fig 3). The involvement of ERK signalling in growth and development necessitates the study to find other components in order to gain more insight into this pathway.

Chemotaxis pathways

Ras regulates chemotaxis by activating various downstream effector proteins (Fig 4). One of the most studied downstream effectors of Ras is PI3K. Ras activates the basal PI3K present at the membrane; through a positive feedback loop via F-actin polymerization, cytosolic PI3K is also recruited at the membrane [56]. *Dictyostelium* has five class I PI3K proteins [111], which all consist of a characteristic Ras-binding domain that specifically interact with GTP-bound Ras proteins [112]. In a chemoattractant gradient PI3K is localized at the leading edge where it catalyzes the production of PIP3 from PI(4,5)P2 while PTEN is localized at the lateral and posterior sides of the cell catalyzing the converse reaction of dephosphorylating PIP3 to PIP2. Due to this reciprocal localization and activity of these two proteins, a PIP3 gradient is formed promoting pseudopod extension at the leading edge whereas PIP2 suppresses lateral pseudopods. *pten*-null cells can sense the direction of gradient and become polarized but they exhibit inefficient chemotaxis with decreased chemotaxis index, speed and directional persistence [113]. Although these initial studies suggested that the PIP3 pathway is essential and acts as a molecular compass for chemotaxis [114,115], recent data suggest that PI3K only plays an important role in directional sensing of shallow gradients [116,117]. Cells with disruptions in all five *pi3k* genes were still capable of performing normal chemotaxis [111,118] in steep gradients, albeit with a reduced speed, indicating the presence of other pathways that mediate directional sensing. A pathway mediating chemotaxis in the absence of PI3K was identified that involved Phospholipase A2 (PLA2) [116,119]. Although not much is known about this pathway apart from the fact that it is regulated by Ca^{2+} uptake and cytosolic Ca^{2+} , however, various studies suggest that PLA2 is redundant for chemotaxis in steep gradients [116,120]. The cells lacking *pi3k* and *pla2* when starved for 7 hours were still able to exhibit chemotaxis suggesting the existence of still some other pathway. Mutational analysis revealed the importance of sGC

pathway in regulating chemotaxis [121]. sGC is localized at the leading edge and is predominantly enriched in the extended pseudopod where it interacts with actin filaments serving as the memory of an existing pseudopod. It also catalyzes the formation of cGMP by hydrolyzing GTP [122]. cGMP induces the formation of myosin filaments thereby also inhibiting the formation of pseudopods at the rear and at lateral sides of the cell [123,124]. RasG regulates the production of cGMP upon cAMP stimulation further signifying the importance of Ras in chemotaxis [81]. Also RasC activates TORC2 which then phosphorylates PKBR1 and PKBA at the leading edge of cell thereby regulating chemotactic responses [93].

Role of Ras in Cytoskeleton Rearrangement

Symmetry breaking in the above described signalling pathways leads to major changes in the cytoskeleton; actin polymerization occurs at the leading edge of the cell, while acto-myosin filaments are formed at the rear and back of the cell. Since a properly organized cytoskeleton is essential for chemotaxis and development, regulation of cytoskeleton is of utmost importance. Actin polymerization at the leading edge is mediated by the Arp2/3 complex which produces a complex network of branched filaments that leads to pseudopod extension. The regulation of this process involves several groups of proteins, including the Rho family of small G-proteins, WASP (Wiskott-Aldrich syndrome protein), and WAVE/SCAR [125-127].

At the lateral and posterior sides of a chemotaxing cell, pseudopod extension is suppressed by assembly of myosin II filaments [128-130] in a cGMP dependent manner [85]. Myosin II is responsible for contraction of rear end as well as involved in spatio-temporal regulation of traction forces [131,132]. Furthermore, at the leading edge activated RapA stimulates Phg2 (a serine/threonine kinase) which phosphorylates and thus leads to disassembly of myosin II [133-135]. Disruption of myosin II leads to inefficient chemotaxis and formation of lateral pseudopodia [136,137].




Figure 4 Signaling pathways regulating chemotaxis Binding of chemoattractant to the receptor leads to the activation of associated heterotrimeric G-protein which further results in rapid activation of Ras at the leading edge. The activation of Ras results in the activation of four signaling pathways along with Rap specific GEFs. The activated PI3K and TorC2 are important for actin polymerization at the leading edge. Rac proteins also play an important role in actin polymerization at the front of the cell. cGMP produced by sGC and GCA is a key regulator of myosin filament formation at the back of the cell. Upon activation, Rap binds to Phg2 and inhibits formation of myosin filaments at the front of the cell. The presence of several positive feedback loops results in stronger Ras activation and F-actin polymerization at the front leading to efficient chemotaxis. The dotted lines indicate pathways that are not completely characterized yet.

Aim of the thesis

Chemotaxis is a complex cellular process involving several interconnected signaling pathways, positive and negative feedback loops, multiple regulators and effector proteins. The information presented in this chapter shows that processes involved in chemotaxis are very complicated. In Chapter 2 of the thesis we show that cells lacking all four known chemotaxis pathways downstream of Ras: sGC, PLA2, PI3K and TORC2, are still able to move. Thus we deduced the basal module for chemotaxis detailing the minimal requirements. Subsequently we characterize Ras activation, symmetry breaking and confinement during *Dictyostelium* chemotaxis using various mutant cell lines in Chapter 3. Nevertheless, in addition to these important components, it is likely that there are many additional proteins, functioning as scaffold, activators, inhibitors and/or effectors in the chemotaxis pathway that still need to be identified. To discover these additional components of the presented basal pathway is a challenging task. Chapter 4 details a new proteomic approach undertaken to find the novel players in chemotactic signal transduction. Ric8, which we found in the screens of several G α proteins, was selected for further studies and shown to be acting as a non-receptor GEF for *Dictyostelium* G α proteins discussed in detail in Chapter 5. Similarly, Chapter 6 deals with characterizing a leucine rich repeat protein LrrA which could be a putative link between G α and Ras signaling.

References

1. Armstrong P.B. (1985). The control of cell motility during embryogenesis. *Cancer Metastasis Rev.* 4: 59-79.
2. Czirok A. and Little C.D. (2012). Pattern formation during vasculogenesis. *Birth Defects Res. C. Embryo. Today* 96: 153-162.
3. Kolaczowska E. and Kubes P. (2013). Neutrophil recruitment and function in health and inflammation. *Nat. Rev. Immunol.* 13: 159-175.
4. Balkwill F. (2004). Cancer and the chemokine network. *Nat. Rev. Cancer* 4: 540-550.
5. Eccles S.A. (2005). Targeting key steps in metastatic tumour progression. *Curr. Opin. Genet. Dev.* 15: 77-86.
6. Drechsler M., Megens R.T., van Z.M., Weber C., and Soehnlein O. (2010). Hyperlipidemia-triggered neutrophilia promotes early atherosclerosis. *Circulation* 122: 1837-1845.
7. Devreotes P.N. and Zigmond S.H. (1988). Chemotaxis in Eukaryotic Cells - A Focus on Leukocytes and *Dictyostelium*. *Annual Review of Cell Biology* 4: 649-686.
8. Eichinger L., Pachebat J.A., Glockner G., Rajandream M.A., Sucgang R., Berriman M., Song J., Olsen R., Szafranski K., Xu Q. et al. (2005). The genome of the social amoeba *Dictyostelium discoideum*. *Nature* 435: 43-57.

9. Fey P., Dodson R.J., Basu S., and Chisholm R.L. (2013). One stop shop for everything Dictyostelium: dictyBase and the Dicty Stock Center in 2012. *Methods Mol. Biol.* 983: 59-92.
10. Parent C.A. (2004). Making all the right moves: chemotaxis in neutrophils and Dictyostelium. *Curr. Opin. Cell Biol.* 16: 4-13.
11. Swaney K.F., Huang C.H., and Devreotes P.N. (2010). Eukaryotic chemotaxis: a network of signaling pathways controls motility, directional sensing, and polarity. *Annu. Rev. Biophys.* 39: 265-289.
12. Devreotes P. and Janetopoulos C. (2003). Eukaryotic chemotaxis: distinctions between directional sensing and polarization. *J. Biol. Chem.* 278: 20445-20448.
13. Van Haastert P.J.M. and Devreotes P.N. (2004). Chemotaxis: Signalling the way forward. *Nature Reviews Molecular Cell Biology* 5: 626-634.
14. Xu J., Wang F., Van K.A., Herzmark P., Straight A., Kelly K., Takuwa Y., Sugimoto N., Mitchison T., and Bourne H.R. (2003). Divergent signals and cytoskeletal assemblies regulate self-organizing polarity in neutrophils. *Cell* 114: 201-214.
15. Levi S., Polyakov M.V., and Egelhoff T.T. (2002). Myosin II dynamics in Dictyostelium: Determinants for filament assembly and translocation to the cell cortex during chemoattractant responses. *Cell Motility and the Cytoskeleton* 53: 177-188.
16. Moores S.L., Sabry J.H., and Spudich J.A. (1996). Myosin dynamics in live Dictyostelium cells. *Proc. Natl. Acad. Sci. U. S. A* 93: 443-446.
17. Parent C.A., Blacklock B.J., Froehlich W.M., Murphy D.B., and Devreotes P.N. (1998). G protein signaling events are activated at the leading edge of chemotactic cells. *Cell* 95: 81-91.
18. Fredriksson R., Lagerstrom M.C., Lundin L.G., and Schioth H.B. (2003). The G-protein-coupled receptors in the human genome form five main families. Phylogenetic analysis, paralogon groups, and fingerprints. *Mol. Pharmacol.* 63: 1256-1272.
19. Prabhu Y. and Eichinger L. (2006). The Dictyostelium repertoire of seven transmembrane domain receptors. *Eur. J. Cell Biol.* 85: 937-946.
20. Louis J.M., Ginsburg G.T., and Kimmel A.R. (1994). The cAMP receptor CAR4 regulates axial patterning and cellular differentiation during late development of Dictyostelium. *Genes Dev.* 8: 2086-2096.
21. Saxe C.L., III, Johnson R., Devreotes P.N., and Kimmel A.R. (1991). Multiple genes for cell surface cAMP receptors in Dictyostelium discoideum. *Dev. Genet.* 12: 6-13.
22. Verkerke-van W., I, Kim J.Y., Brandt R., Devreotes P.N., and Schaap P. (1998). Functional promiscuity of gene regulation by serpentine receptors in Dictyostelium discoideum. *Mol. Cell Biol.* 18: 5744-5749.
23. Sun T.J. and Devreotes P.N. (1991). Gene Targeting of the Aggregation Stage Camp Receptor Car1 in Dictyostelium. *Genes & Development* 5: 572-582.
24. Saxe C.L., III, Johnson R.L., Devreotes P.N., and Kimmel A.R. (1991). Expression of a cAMP receptor gene of Dictyostelium and evidence for a multigene family. *Genes Dev.* 5: 1-8.
25. Yu Y. and Saxe C.L., III (1996). Differential distribution of cAMP receptors cAR2 and cAR3 during Dictyostelium development. *Dev. Biol.* 173: 353-356.
26. Saxe C.L., III, Ginsburg G.T., Louis J.M., Johnson R., Devreotes P.N., and Kimmel A.R. (1993). CAR2, a prestalk cAMP receptor required for normal tip formation and late development of Dictyostelium discoideum. *Genes Dev.* 7: 262-272.
27. Johnson R.L., Saxe C.L., III, Gollop R., Kimmel A.R., and Devreotes P.N. (1993). Identification and targeted gene disruption of cAR3, a cAMP receptor subtype expressed during multicellular stages of Dictyostelium development. *Genes Dev.* 7: 273-282.

28. Coleman D.E., Berghuis A.M., Lee E., Linder M.E., Gilman A.G., and Sprang S.R. (1994). Structures of active conformations of Gi alpha 1 and the mechanism of GTP hydrolysis. *Science* 265: 1405-1412.
29. Oldham W.M. and Hamm H.E. (2008). Heterotrimeric G protein activation by G-protein-coupled receptors. *Nat. Rev. Mol. Cell Biol.* 9: 60-71.
30. Bornancin F., Pfister C., and Chabre M. (1989). The transitory complex between photoexcited rhodopsin and transducin. Reciprocal interaction between the retinal site in rhodopsin and the nucleotide site in transducin. *Eur. J. Biochem.* 184: 687-698.
31. Koelle M.R. (1997). A new family of G-protein regulators - the RGS proteins. *Curr. Opin. Cell Biol.* 9: 143-147.
32. Hepler J.R. (1999). Emerging roles for RGS proteins in cell signalling. *Trends Pharmacol. Sci.* 20: 376-382.
33. Clapham D.E. and Neer E.J. (1997). G protein beta gamma subunits. *Annual Review of Pharmacology and Toxicology* 37: 167-203.
34. Ford C.E., Skiba N.P., Bae H.S., Daaka Y.H., Reuveny E., Shekter L.R., Rosal R., Weng G.Z., Yang C.S., Iyengar R. et al. (1998). Molecular basis for interactions of G protein beta gamma subunits with effectors. *Science* 280: 1271-1274.
35. Lilly P., Wu L., Welker D.L., and Devreotes P.N. (1993). A G-protein beta-subunit is essential for Dictyostelium development. *Genes Dev.* 7: 986-995.
36. Zhang N., Long Y., and Devreotes P.N. (2001). Ggamma in dictyostelium: its role in localization of gbetagamma to the membrane is required for chemotaxis in shallow gradients. *Mol. Biol. Cell* 12: 3204-3213.
37. Li Y., Sternweis P.M., Charnecki S., Smith T.F., Gilman A.G., Neer E.J., and Kozasa T. (1998). Sites for Galpha binding on the G protein beta subunit overlap with sites for regulation of phospholipase Cbeta and adenylyl cyclase. *J. Biol. Chem.* 273: 16265-16272.
38. Jin T., Amzel M., Devreotes P.N., and Wu L. (1998). Selection of gbeta subunits with point mutations that fail to activate specific signaling pathways in vivo: dissecting cellular responses mediated by a heterotrimeric G protein in Dictyostelium discoideum. *Mol. Biol. Cell* 9: 2949-2961.
39. Brzustowski J.A., Johnson C., and Kimmel A.R. (2002). Galpha-mediated inhibition of developmental signal response. *Curr. Biol.* 12: 1199-1208.
40. Kumagai A., Pupillo M., Gundersen R., Miakelye R., Devreotes P.N., and Firtel R.A. (1989). Regulation and Function of G-Alpha Protein Subunits in Dictyostelium. *Cell* 57: 265-275.
41. Kumagai A., Hadwiger J.A., Pupillo M., and Firtel R.A. (1991). Molecular Genetic-Analysis of 2 G-Alpha Protein Subunits in Dictyostelium. *Journal of Biological Chemistry* 266: 1220-1228.
42. Hadwiger J.A., Natarajan K., and Firtel R.A. (1996). Mutations in the Dictyostelium heterotrimeric G protein alpha subunit G alpha5 alter the kinetics of tip morphogenesis. *Development* 122: 1215-1224.
43. Wu L.J., Valkema R., Vanhaastert P.J.M., and Devreotes P.N. (1995). The G-Protein Beta-Subunit Is Essential for Multiple Responses to Chemoattractants in Dictyostelium. *Journal of Cell Biology* 129: 1667-1675.
44. Brandon M.A. and Podgorski G.J. (1997). G alpha 3 regulates the cAMP signaling system in Dictyostelium. *Mol. Biol. Cell* 8: 1677-1685.
45. Hadwiger J.A., Lee S., and Firtel R.A. (1994). The G(Alpha) Subunit G(Alpha)4 Couples to Pterin Receptors and Identifies A Signaling Pathway That Is Essential for Multicellular Development in Dictyostelium. *Proceedings of the National Academy of Sciences of the United States of America* 91: 10566-10570.

46. Okaichi K., Cubitt A.B., Pitt G.S., and Firtel R.A. (1992). Amino-Acid Substitutions in the Dictyostelium G-Alpha Subunit G-Alpha-2 Produce Dominant Negative Phenotypes and Inhibit the Activation of Adenylyl Cyclase, Guanylyl Cyclase, and Phospholipase-C. *Molecular Biology of the Cell* 3: 735-747.
47. Hadwiger J.A. and Firtel R.A. (1992). Analysis of G alpha 4, a G-protein subunit required for multicellular development in Dictyostelium. *Genes Dev.* 6: 38-49.
48. Hadwiger J.A. and Srinivasan J. (1999). Folic acid stimulation of the Galpha4 G protein-mediated signal transduction pathway inhibits anterior prestalk cell development in Dictyostelium. *Differentiation* 64: 195-204.
49. Jin T., Zhang N., Long Y., Parent C.A., and Devreotes P.N. (2000). Localization of the G protein beta gamma complex in living cells during chemotaxis. *Science* 287: 1034-1036.
50. Janetopoulos C., Jin T., and Devreotes P. (2001). Receptor-mediated activation of heterotrimeric G-proteins in living cells. *Science* 291: 2408-2411.
51. Xiao Z., Zhang N., Murphy D.B., and Devreotes P.N. (1997). Dynamic distribution of chemoattractant receptors in living cells during chemotaxis and persistent stimulation. *Journal of Cell Biology* 139: 365-374.
52. Ueda M., Sako Y., Tanaka T., Devreotes P., and Yanagida T. (2001). Single-molecule analysis of chemotactic signaling in Dictyostelium cells. *Science* 294: 864-867.
53. Janetopoulos C., Ma L., Devreotes P.N., and Iglesias P.A. (2004). Chemoattractant-induced phosphatidylinositol 3,4,5-trisphosphate accumulation is spatially amplified and adapts, independent of the actin cytoskeleton. *Proc. Natl. Acad. Sci. U. S. A* 101: 8951-8956.
54. Xu X., Meier-Schellersheim M., Jiao X., Nelson L.E., and Jin T. (2005). Quantitative imaging of single live cells reveals spatiotemporal dynamics of multistep signaling events of chemoattractant gradient sensing in Dictyostelium. *Mol. Biol. Cell* 16: 676-688.
55. Kae H., Kortholt A., Rehmann H., Insall R.H., Van Haastert P.J., Spiegelman G.B., and Weeks G. (2007). Cyclic AMP signalling in Dictyostelium: G-proteins activate separate Ras pathways using specific RasGEFs. *EMBO Rep.* 8: 477-482.
56. Sasaki A.T., Chun C., Takeda K., and Firtel R.A. (2004). Localized Ras signaling at the leading edge regulates PI3K, cell polarity, and directional cell movement. *J. Cell Biol.* 167: 505-518.
57. Dharmawardhane S., Cubitt A.B., Clark A.M., and Firtel R.A. (1994). Regulatory role of the G alpha 1 subunit in controlling cellular morphogenesis in Dictyostelium. *Development* 120: 3549-3561.
58. Kumagai A., Pupillo M., Gundersen R., Miakelye R., Devreotes P.N., and Firtel R.A. (1989). Regulation and Function of G-Alpha Protein Subunits in Dictyostelium. *Cell* 57: 265-275.
59. Kumagai A., Hadwiger J.A., Pupillo M., and Firtel R.A. (1991). Molecular Genetic-Analysis of 2 G-Alpha Protein Subunits in Dictyostelium. *Journal of Biological Chemistry* 266: 1220-1228.
60. Brandon M.A., Voglmaier S., and Siddiqi A.A. (1997). Molecular characterization of a Dictyostelium G-protein alpha-subunit required for development. *Gene* 200: 99-105.
61. Brandon M.A., Mahadeo D.C., and Podgorski G.J. (2002). Galpha3 and protein kinase A represent cross-talking pathways for gene expression in Dictyostelium discoideum. *Dev. Growth Differ.* 44: 457-465.
62. Srinivasan J., Gundersen R.E., and Hadwiger J.A. (1999). Activated Galpha subunits can inhibit multiple signal transduction pathways during Dictyostelium development. *Dev. Biol.* 215: 443-452.
63. Nguyen H.N., Raisley B., and Hadwiger J.A. (2010). MAP kinases have different functions in Dictyostelium G protein-mediated signaling. *Cell Signal.* 22: 836-847.
64. Raisley B., Nguyen H.N., and Hadwiger J.A. (2010). G{alpha}5 subunit-mediated signalling requires a D-motif and the MAPK ERK1 in Dictyostelium. *Microbiology* 156: 789-797.

65. Wu L., Gaskins C., Zhou K., Firtel R.A., and Devreotes P.N. (1994). Cloning and targeted mutations of G alpha 7 and G alpha 8, two developmentally regulated G protein alpha-subunit genes in Dictyostelium. *Mol. Biol. Cell* 5: 691-702.
66. Wu Y. and Janetopoulos C. (2013). The G alpha subunit Galpha8 inhibits proliferation, promotes adhesion and regulates cell differentiation. *Dev. Biol.* 380: 58-72.
67. Bakthavatsalam D., Choe J.M., Hanson N.E., and Gomer R.H. (2009). A Dictyostelium chalone uses G proteins to regulate proliferation. *BMC. Biol.* 7: 44.
68. Phillips J.E. and Gomer R.H. (2012). A secreted protein is an endogenous chemorepellant in Dictyostelium discoideum. *Proc. Natl. Acad. Sci. U. S. A* 109: 10990-10995.
69. Kortholt A. and Van Haastert P.J. (2008). Highlighting the role of Ras and Rap during Dictyostelium chemotaxis. *Cell Signal*.
70. Weeks G. (2005). The small GTPase superfamily. In Dictyostelium genomics, William F.Loomis, Adam Kuspa, eds. Horizon Bioscience), pp. 173-210.
71. Daniel J., Spiegelman G.B., and Weeks G. (1993). Characterization of a third ras gene, rasB, that is expressed throughout the growth and development of Dictyostelium discoideum. *Oncogene* 8: 1041-1047.
72. Sutherland B.W., Spiegelman G.B., and Weeks G. (2001). A Ras subfamily GTPase shows cell cycle-dependent nuclear localization. *Embo Reports* 2: 1024-1028.
73. Lim C.J., Spiegelman G.B., and Weeks G. (2001). RasC is required for optimal activation of adenylyl cyclase and Akt/PKB during aggregation. *Embo Journal* 20: 4490-4499.
74. Lim C.J., Zawadzki K.A., Khosla M., Secko D.M., Spiegelman G.B., and Weeks G. (2005). Loss of the Dictyostelium RasC protein alters vegetative cell size, motility and endocytosis. *Experimental Cell Research* 306: 47-55.
75. Wessels D., Brincks R., Kuhl S., Stepanovic V., Daniels K.J., Weeks G., Lim C.J., Spiegelman G., Fuller D., Iranfar N. et al. (2004). RasC plays a role in transduction of temporal gradient information in the cyclic-AMP wave of Dictyostelium discoideum. *Eukaryot. Cell* 3: 646-662.
76. Kae H., Lim C.J., Spiegelman G.B., and Weeks G. (2004). Chemoattractant-induced Ras activation during Dictyostelium aggregation. *Embo Reports* 5: 602-606.
77. Raymond C.D., Gomer R.H., Mehdy M.C., and Firtel R.A. (1984). Developmental Regulation of A Dictyostelium Gene Encoding A Protein Homologous to Mammalian Ras Protein. *Cell* 39: 141-148.
78. Raymond C.D., Gomer R.H., Nellen W., Theibert A., Devreotes P., and Firtel R.A. (1986). Phenotypic changes induced by a mutated ras gene during the development of Dictyostelium transformants. *Nature* 323: 340-343.
79. Khosla M., Robbins S.M., Spiegelman G.B., and Weeks G. (1990). Regulation of DdRasG gene expression during Dictyostelium development. *Mol. Cell Biol.* 10: 918-922.
80. Tuxworth R.I., Cheetham J.L., Machesky L.M., Spiegelmann G.B., Weeks G., and Insall R.H. (1997). Dictyostelium RasG is required for normal motility and cytokinesis, but not growth. *Journal of Cell Biology* 138: 605-614.
81. Bolourani P., Spiegelman G.B., and Weeks G. (2006). Delineation of the roles played by RasG and RasC in cAMP-dependent signal transduction during the early development of Dictyostelium discoideum. *Mol. Biol. Cell* 17: 4543-4550.
82. Chubb J.R., Wilkins A., Thomas G.M., and Insall R.H. (2000). The Dictyostelium RasS protein is required for macropinocytosis, phagocytosis and the control of cell movement. *Journal of Cell Science* 113: 709-719.
83. Chubb J.R., Wilkins A., Wessels D.J., Soll D.R., and Insall R.H. (2002). Pseudopodium dynamics and rapid cell movement in Dictyostelium Ras pathway mutants. *Cell Motility and the Cytoskeleton* 53: 150-162.

84. Daniel J., Bush J., Cardelli J., Spiegelman G.B., and Weeks G. (1994). Isolation of two novel ras genes in *Dictyostelium discoideum*; evidence for a complex, developmentally regulated ras gene subfamily. *Oncogene* 9: 501-508.
85. Jeon T.J., Lee D.J., Merlot S., Weeks G., and Firtel R.A. (2007). Rap1 controls cell adhesion and cell motility through the regulation of myosin II. *Journal of Cell Biology* 176: 1021-1033.
86. Jeon T.J., Lee D.J., Lee S., Weeks G., and Firtel R.A. (2007). Regulation of Rap1 activity by RapGAP1 controls cell adhesion at the front of chemotaxing cells. *J. Cell Biol.* 179: 833-843.
87. Kang R.J., Kae H., Ip H., Spiegelman G.B., and Weeks G. (2002). Evidence for a role for the *Dictyostelium* Rap1 in cell viability and the response to osmotic stress. *Journal of Cell Science* 115: 3675-3682.
88. Robbins S.M., Suttrop V.V., Weeks G., and Spiegelman G.B. (1990). A ras-related gene from the lower eukaryote *Dictyostelium* that is highly conserved relative to the human rap genes. *Nucleic Acids Res.* 18: 5265-5269.
89. Wilkins A., Szafranski K., Fraser D.J., Bakthavatsalam D., Muller R., Fisher P.R., Glockner G., Eichinger L., Noegel A.A., and Insall R.H. (2005). The *Dictyostelium* genome encodes numerous RasGEFs with multiple biological roles. *Genome Biol.* 6: R68.
90. Zhang S., Charest P.G., and Firtel R.A. (2008). Spatiotemporal regulation of Ras activity provides directional sensing. *Curr. Biol.* 18: 1587-1593.
91. Insall R.H., Borleis J., and Devreotes P.N. (1996). The aimless RasGEF is required for processing of chemotactic signals through G-protein-coupled receptors in *Dictyostelium*. *Current Biology* 6: 719-729.
92. Pitt G.S., Milona N., Borleis J., Lin K.C., Reed R.R., and Devreotes P.N. (1992). Structurally Distinct and Stage-Specific Adenylyl Cyclase Genes Play Different Roles in *Dictyostelium* Development. *Cell* 69: 305-315.
93. Cai H., Das S., Kamimura Y., Long Y., Parent C.A., and Devreotes P.N. (2010). Ras-mediated activation of the TORC2-PKB pathway is critical for chemotaxis. *J. Cell Biol.* 190: 233-245.
94. Charest P.G., Shen Z., Lakoduk A., Sasaki A.T., Briggs S.P., and Firtel R.A. (2010). A Ras signaling complex controls the RasC-TORC2 pathway and directed cell migration. *Dev. Cell* 18: 737-749.
95. Lee S., Comer F.I., Sasaki A., McLeod I.X., Duong Y., Okumura K., Yates J.R., III, Parent C.A., and Firtel R.A. (2005). TOR complex 2 integrates cell movement during chemotaxis and signal relay in *Dictyostelium*. *Mol. Biol. Cell* 16: 4572-4583.
96. Lee S., Parent C.A., Insall R., and Firtel R.A. (1999). A novel Ras-interacting protein required for chemotaxis and cyclic adenosine monophosphate signal relay in *Dictyostelium*. *Mol. Biol. Cell* 10: 2829-2845.
97. Chen M.Y., Long Y., and Devreotes P.N. (1997). A novel cytosolic regulator, pianissimo, is required for chemoattractant receptor and G protein-mediated activation of the 12 transmembrane domain adenylyl cyclase in *Dictyostelium*. *Genes & Development* 11: 3218-3231.
98. Maeda M., Aubry L., Insall R., Gaskins C., Devreotes P.N., and Firtel R.A. (1996). Seven helix chemoattractant receptors transiently stimulate mitogen-activated protein kinase in *Dictyostelium*. Role of heterotrimeric G proteins. *J. Biol. Chem.* 271: 3351-3354.
99. Segall J.E., Kuspa A., Shauly G., Ecke M., Maeda M., Gaskins C., Firtel R.A., and Loomis W.F. (1995). A MAP kinase necessary for receptor-mediated activation of adenylyl cyclase in *Dictyostelium*. *J. Cell Biol.* 128: 405-413.
100. Gaskins C., Maeda M., and Firtel R.A. (1994). Identification and functional analysis of a developmentally regulated extracellular signal-regulated kinase gene in *Dictyostelium discoideum*. *Mol. Cell Biol.* 14: 6996-7012.

101. Lewis T.S., Shapiro P.S., and Ahn N.G. (1998). Signal transduction through MAP kinase cascades. *Adv. Cancer Res.* 74: 49-139.
102. Pearson G., Robinson F., Beers G.T., Xu B.E., Karandikar M., Berman K., and Cobb M.H. (2001). Mitogen-activated protein (MAP) kinase pathways: regulation and physiological functions. *Endocr. Rev.* 22: 153-183.
103. Chang L. and Karin M. (2001). Mammalian MAP kinase signalling cascades. *Nature* 410: 37-40.
104. Sommers C.L., Gurson J.M., Surana R., Barda-Saad M., Lee J., Kishor A., Li W., Gasser A.J., Barr V.A., Miyaji M. et al. (2008). Bam32: a novel mediator of Erk activation in T cells. *Int. Immunol.* 20: 811-818.
105. Kortum R.L., Rouquette-Jazdanian A.K., and Samelson L.E. (2013). Ras and extracellular signal-regulated kinase signaling in thymocytes and T cells. *Trends Immunol.* 34: 259-268.
106. Ma H., Gamper M., Parent C., and Firtel R.A. (1997). The Dictyostelium MAP kinase kinase DdMEK1 regulates chemotaxis and is essential for chemoattractant-mediated activation of guanylyl cyclase. *EMBO J.* 16: 4317-4332.
107. Chung C.Y., Reddy T.B., Zhou K., and Firtel R.A. (1998). A novel, putative MEK kinase controls developmental timing and spatial patterning in Dictyostelium and is regulated by ubiquitin-mediated protein degradation. *Genes Dev.* 12: 3564-3578.
108. Gaskins C., Clark A.M., Aubry L., Segall J.E., and Firtel R.A. (1996). The Dictyostelium MAP kinase ERK2 regulates multiple, independent developmental pathways. *Genes Dev.* 10: 118-128.
109. Wang Y., Liu J., and Segall J.E. (1998). MAP kinase function in amoeboid chemotaxis. *J. Cell Sci.* 111 (Pt 3): 373-383.
110. Maeda M., Lu S., Shaulsky G., Miyazaki Y., Kuwayama H., Tanaka Y., Kuspa A., and Loomis W.F. (2004). Periodic signaling controlled by an oscillatory circuit that includes protein kinases ERK2 and PKA. *Science* 304: 875-878.
111. Hoeller O. and Kay R.R. (2007). Chemotaxis in the absence of PIP3 gradients. *Curr. Biol.* 17: 813-817.
112. Herrmann C. (2003). Ras-effector interactions: after one decade. *Curr. Opin. Struct. Biol.* 13: 122-129.
113. Wessels D., Lusche D.F., Kuhl S., Heid P., and Soll D.R. (2007). PTEN plays a role in the suppression of lateral pseudopod formation during Dictyostelium motility and chemotaxis. *J. Cell Sci.* 120: 2517-2531.
114. Funamoto S., Milan K., Meili R., and Firtel R.A. (2001). Role of phosphatidylinositol 3' kinase and a downstream pleckstrin homology domain-containing protein in controlling chemotaxis in Dictyostelium. *Journal of Cell Biology* 153: 795-809.
115. Funamoto S., Meili R., Lee S., Parry L., and Firtel R.A. (2002). Spatial and temporal regulation of 3-phosphoinositides by PI 3-kinase and PTEN mediates chemotaxis. *Cell* 109: 611-623.
116. van Haastert P.J., Keizer-Gunnink I., and Kortholt A. (2007). Essential role of PI3-kinase and phospholipase A2 in Dictyostelium discoideum chemotaxis. *J. Cell Biol.* 177: 809-816.
117. Takeda K., Sasaki A.T., Ha H.J., Seung H.A., and Firtel R.A. (2007). Role of phosphatidylinositol 3-kinases in chemotaxis in Dictyostelium. *Journal of Biological Chemistry* 282: 11874-11884.
118. Van Haastert P.J. and Veltman D.M. (2007). Chemotaxis: navigating by multiple signaling pathways. *Sci. STKE.* 2007: e40.
119. Chen L., Iijima M., Tang M., Landree M.A., Huang Y.E., Xiong Y., Iglesias P.A., and Devreotes P.N. (2007). PLA(2) and PI3K/PTEN Pathways Act in Parallel to Mediate Chemotaxis. *Dev. Cell* 12: 603-614.
120. Kortholt A., Kataria R., Keizer-Gunnink I., van Egmond W.N., Khanna A., and Van Haastert P.J. (2011). Dictyostelium chemotaxis: essential Ras activation and accessory signalling pathways for amplification. *EMBO Rep.* 12: 1273-1279.

121. Veltman D.M., Keizer-Gunnink I., and Van Haastert P.J. (2008). Four key signaling pathways mediating chemotaxis in *Dictyostelium discoideum*. *J. Cell Biol.* **180**: 747-753.
122. Roelofs J. and Van Haastert P.J. (2002). Characterization of two unusual guanylyl cyclases from *dictyostelium*. *J. Biol. Chem.* **277**: 9167-9174.
123. Bosgraaf L., Russcher H., Smith J.L., Wessels D., Soll D.R., and Van Haastert P.J.M. (2002). A novel cGMP signalling pathway mediating myosin phosphorylation and chemotaxis in *Dictyostelium*. *Embo Journal* **21**: 4560-4570.
124. Veltman D.M. and Van Haastert P.J. (2006). Guanylyl cyclase protein and cGMP product independently control front and back of chemotaxing *Dictyostelium* cells. *Mol. Biol. Cell* **17**: 3921-3929.
125. Ibarra N., Pollitt A., and Insall R.H. (2005). Regulation of actin assembly by SCAR/WAVE proteins. *Biochem. Soc. Trans.* **33**: 1243-1246.
126. Machesky L.M. and Insall R.H. (1998). Scar1 and the related Wiskott-Aldrich syndrome protein, WASP, regulate the actin cytoskeleton through the Arp2/3 complex. *Curr. Biol.* **8**: 1347-1356.
127. Pollitt A.Y. and Insall R.H. (2009). WASP and SCAR/WAVE proteins: the drivers of actin assembly. *J. Cell Sci.* **122**: 2575-2578.
128. Chung C.Y., Potikyan G., and Firtel R.A. (2001). Control of cell polarity and chemotaxis by Akt/PKB and PI3 kinase through the regulation of PAKs. *Molecular Cell* **7**: 937-947.
129. Ridley A.J., Schwartz M.A., Burridge K., Firtel R.A., Ginsberg M.H., Borisy G., Parsons J.T., and Horwitz A.R. (2003). Cell migration: integrating signals from front to back. *Science* **302**: 1704-1709.
130. Stephens L., Milne L., and Hawkins P. (2008). Moving towards a better understanding of chemotaxis. *Curr. Biol.* **18**: R485-R494.
131. Reichl E.M., Ren Y., Morphew M.K., Delannoy M., Effler J.C., Girard K.D., Divi S., Iglesias P.A., Kuo S.C., and Robinson D.N. (2008). Interactions between myosin and actin crosslinkers control cytokinesis contractility dynamics and mechanics. *Curr. Biol.* **18**: 471-480.
132. Meili R., Alonso-Latorre B., del Alamo J.C., Firtel R.A., and Lasheras J.C. (2010). Myosin II is essential for the spatiotemporal organization of traction forces during cell motility. *Mol. Biol. Cell* **21**: 405-417.
133. Rebstein P.J., Cardelli J., Weeks G., and Spiegelman G.B. (1997). Mutational analysis of the role of Rap1 in regulating cytoskeletal function in *Dictyostelium*. *Experimental Cell Research* **231**: 276-283.
134. Gebbie L., Benghezal M., Cornillon S., Froquet R., Cherix N., Malbouyres M., Lefkir Y., Grangeasse C., Fache S., Dalous J. et al. (2004). Phg2, a kinase involved in adhesion and focal site modeling in *Dictyostelium*. *Mol. Biol. Cell* **15**: 3915-3925.
135. Kortholt A., Rehmann H., Kae H., Bosgraaf L., Keizer-Gunnink I., Weeks G., Wittinghofer A., and Van Haastert P.J. (2006). Characterization of the GbpD-activated Rap1 pathway regulating adhesion and cell polarity in *Dictyostelium discoideum*. *J. Biol. Chem.* **281**: 23367-23376.
136. Bosgraaf L. and van Haastert P.J. (2006). The regulation of myosin II in *Dictyostelium*. *Eur. J. Cell Biol.* **85**: 969-979.
137. Yumura S., Yoshida M., Betapudi V., Licate L.S., Iwadata Y., Nagasaki A., Uyeda T.Q., and Egelhoff T.T. (2005). Multiple myosin II heavy chain kinases: roles in filament assembly control and proper cytokinesis in *Dictyostelium*. *Mol. Biol. Cell* **16**: 4256-4266.